

נספח 2. מקורות

- [1] כבביה ד. 1994. התנהגות רעיה והזנה של עיזים במרעה חורש ים-תיכוני (עבודת גמר לתואר מוסמך). רחובות: האוניברסיטה העברית בירושלים.
- [2] נבון ש. 2014. מבט מקרוב על התנהגות הרעייה של עיזים בחורש ים תיכוני (עבודת גמר לתואר מוסמך). רחובות: האוניברסיטה העברית בירושלים.
- [3] נבון ש, קיגל ח, חודאי נואנגר י"ד. 2016. שונות כימית באוכלוסייה מקומית של אלת המסטיק (*Pistacia lentiscus* L.). יער 16: 36-45.
- [4] פרבולוצקי א. 1991. נקמת השעיר לעזאזל. טבע וארץ ל"ג: 4.
- [5] שביב ע. 1978. אוטאקולוגיה של אלת המסטיק (*Pistacia lentiscus* L.) (עבודה לקבלת תואר דוקטור). חיפה: הטכניון – מכון טכנולוגי לישראל.
- [6] שמידע א. 1985. עושר המינים והאבולוציה של המינים החד-שנתיים בחורש הים-תיכוני. עלון רתם 18: 57-68.
- [7] Arviv A, Muklada H, Kigel J, et al. 2016. Targeted grazing of milk thistle (*Silybum marianum*) and Syrian thistle (*Notobasis syriaca*) by goats: Preference following preconditioning, generational transfer, and toxicity. *Applied Animal Behaviour Science* 179: 53-59.
- [8] Avrani S, Ben-Shlomo R, and Inbar M. 2012. Genetic structure of a galling aphid *Slavum wertheimae* and its host tree *Pistacia atlantica* across an Irano-Turanian distribution: From fragmentation to speciation? *Tree Genetics & Genomes* 8: 811-820.
- [9] Baraza E, Hódar JA, and Zamora R. 2009. Consequences of plant–chemical diversity for domestic goat food preference in Mediterranean forests. *Acta Oecologica* 35: 117-127.
- [10] Bashan D and Bar-Massada A. 2017. Regeneration dynamics of woody vegetation in a Mediterranean landscape under different disturbance-based management treatments. *Applied Vegetation Science* 20: 106-114.
- [11] Bedoya-Pérez MA, Isler I, Banks PB, and McArthur C. 2014. Roles of the volatile terpene, 1, 8-cineole, in plant–herbivore interactions: A foraging odor cue as well as a toxin?. *Oecologia* 174: 827-837.

- [12] Decandia M, Sitzia M, Cabiddu A, et al. 2000. The use of polyethylene glycol to reduce the anti-nutritional effects of tannins in goats fed woody species. *Small Ruminant Research* **38**: 157-164.
- [13] Delphia CM, Rohr JR, Stephenson AG, et al. 2009. Effects of genetic variation and inbreeding on volatile production in a field population of horsenettle. *International Journal of Plant Sciences* **170**: 12-20.
- [14] Dicke M and Baldwin IT. 2010. The evolutionary context for herbivore-induced plant volatiles: Beyond the 'cry for help'. *Trends in Plant Science* **15**: 167-175.
- [15] Hamrouni-Aschi E, Larbi Khouja M, Boussaid M, et al. 2013. Essential oil composition of the Tunisian endemic cypress (*Cupressus sempervirens* L. var. *numidica* Trab.). *Chemistry & Biodiversity* **10**: 989-1003.
- [16] Estell RE, Fredrickson EL, Tellez MR, et al. 1998. Effects of volatile compounds on consumption of alfalfa pellets by sheep. *Journal of Animal Science* **76**: 228-233.
- [17] Evlagon D, Kommisarchik S, Glasser T, et al. 2010. How much browse is available for goats that graze Mediterranean woodlands?. *Small Ruminant Research* **94**: 103-108.
- [18] Glasser T, Landau S, Ungar ED, et al. 2008. A fecal near-infrared reflectance spectroscopy-aided methodology to determine goat dietary composition in a Mediterranean shrubland. *Journal of Animal Science* **86**: 1345-1356.
- [19] Glasser TA, Landau SY, Ungar ED, et al. 2012. Foraging selectivity of three goat breeds in a Mediterranean shrubland. *Small Ruminant Research* **102**: 7-12.
- [20] Goetsch AL, Gipson TA, Askar AR, and Puchala R. 2010. Feeding behavior of goats. *Journal of Animal Science* **88**: 361-373.
- [21] Goff SA and Klee HJ. 2006. Plant volatile compounds: Sensory cues for health and nutritional value? *Science* **311**: 815-819.

- [22] Korol L, Shklar G, and Schiller G. 2004. Tabor oak in Israel, genetic diversity within and between populations. *Forest Genetics* **11**: 317-323.
- [23] Landau S, Muklada H, Markovics A, and Azaizeh H. 2014. Traditional uses of *Pistacia lentiscus* in veterinary and human medicine. In: Yaniv Z and Dudai N (Eds). *Medicinal and Aromatic Plants of the Middle-East*. Netherlands: Springer.
- [24] Llusia J and Penuelas J. 1998. Changes in terpene content and emission in potted Mediterranean woody plants under severe drought. *Canadian Journal of Botany* **76**: 1366-1373.
- [25] Macheboeuf D, Morgavi DP, Papon Y, et al. 2008. Dose-response effects of essential oils on in vitro fermentation activity of the rumen microbial population. *Animal Feed Science Technology* **145**: 335-350.
- [26] Makkar HPS. 2003. Effect and fate of tannins in ruminant animals, adaptation to tannins and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Ruminant Research* **49**: 241-256.
- [27] Massei G, Cotterill JV, Coats JC, et al. 2007. Can Batesian mimicry help plants to deter herbivores? *Pest Management Science* **63**: 559-563.
- [28] Moore BD, Wallis IR, Palá-Paúl J, et al. 2004. Antiherbivore chemistry of Eucalyptus – Cues and deterrents for marsupial folivores. *Journal of Chemical Ecology* **30**: 1743-1769.
- [29] Nahum S, Inbar M, Ne'eman G, and Ben-Shlomo R. 2008. Phenotypic plasticity and gene diversity in *Pistacia lentiscus* L. along environmental gradients in Israel. *Tree Genetics and Genomes* **4**: 777-785.
- [30] Nagy JG and Tengerdy RP. 1968. Antibacterial action of essential oils of *Artemisia* as an ecological factor II. *Applied Microbiology* **16**: 441-444.
- [31] Ne'eman G. 1993. Variation in leaf phenology and habit in *Quercus ithaburensis*, a Mediterranean deciduous tree. *Journal of Ecology* **81**: 627-634.
- [32] Papachristou TG, Dziba LE, and Provenza FD. 2005. Foraging ecology of goats and sheep on wooded rangelands. *Small Ruminant Research* **59**: 141-156.

- [33] Peñuelas J and Llusà J. 2004. Plant VOC emissions: Making use of the unavoidable. *Trends in Ecology and Evolution* **19**: 402-404.
- [34] Perevolotsky A and Seligman NAG. 1998. Role of grazing in Mediterranean rangeland ecosystems. *BioScience* **48**: 1007-1017.
- [35] Provenza FD, Villalba JJ, Dziba LE, et al. 2003. Linking herbivore experience, varied diets and plant biochemistry diversity. *Small Ruminant Research* **49**: 257-274.
- [36] Riddle RR, Taylor Jr CA, Kothmann MM, and Huston JE. 1996. Volatile oil contents of ashe and redberry juniper and its relationship to preference by Angora and Spanish goats. *Journal of Range Management* **49**: 35-41.
- [37] Rogosic J, Pfister JA, Provenza FD, and Grbesa D. 2006. Sheep and goat preference for and nutritional value of Mediterranean maquis shrubs. *Small Ruminant Research* **64**: 169-179.
- [38] Schiller G, Herr N, Shklar G, and Korol L. 2005. Short Communication: Diversity assessment of leaf phenology variation in *Quercus ithaburensis* Decne by RAPD. *Israel Journal of Plant Sciences* **53**: 75-78.
- [39] Stutz RS, Banks PB, Proschogo N, and McArthur C. 2016. Follow your nose: leaf odour as an important foraging cue for mammalian herbivores. *Oecologia* **182**: 643-651.
- [40] Utsumi SA, Cibils AF, Estell RE, et al. 2009. Seasonal change in one seed juniper intake by sheep and goats in relation to dietary protein and plant secondary metabolites. *Small Ruminant Research* **81**: 152-162.
- [41] Villalba JJ, Provenza FD, and Bryant JP. 2002. Consequences of the interaction between nutrient and plant secondary metabolites on herbivore selectivity: Benefits or detriments for plants? *Oikos* **97**: 282-292.
- [42] Villalba JJ, Provenza FD, and Olson KC. 2006. Terpenes and carbohydrate source influence rumen fermentation, digestibility, intake and preference in sheep. *Journal of Animal Science* **84**: 2463-2473.